

XIII. *Occultatio Palilicii, A. 1738. d. 23. Decembris St. N. observata a D. Christfried Kirchio, Astronomo Regio, Berolini. Ex literis ad J. F. Weidlerum datis excerpta.*

Tempus horolog. oscillar. H. M. S.		Partes microm. Tubo l. ped	Valor partium microm. ' "	Tempus correc- tum. H. M. S.	Tempus correc- tius. H. M. S.
1 56 37	* a centro <i>M. Sinai</i>	38	15 21	6 24 37	
2 58 30	* a centro <i>M. Sinai</i>	34	13 42	26 30	
3 7 3 54	Immersio stellæ			6 31 54	6 32 00
4 8 5 29	Emergio stellæ tub. 9 ped			7 33 29	7 33 33
5 5 35	Emergio certo facta			7 33 35	
6 6 43	* a centro <i>M. Sinai</i>	36	14 32	34 43	
7 15 32	* a prox. marg. Lunæ	8	3 14	43 32	
8 20	* a <i>M. Sinai</i>	46½	18 47	48	
9 28	Diameter Lunæ	74	29 73	56	
	vel	73½	29 41		
10 29 12	<i>M. Sinai</i> a prox. marg. ☉	8	3 14	7 57 12	
11 38	Diameter ☽ sub. 9 ped.	99½	29 38	8 6	
12 42	Diameter ☽ sub. 7 ped.	75	30	10	
13 46 55	* a <i>M. Sinai</i>	72	29 5	14 55	
14 8 48	Diameter Lunæ	73½	29 41	8 16	

Vide SCHEMA hujus Occultationis in TAB. II. Fig. 2.

*Situs stellæ respectu macularum lunarium sequenti modo observatus.*

I. Ante observationem (1.) h. 6. 20'. notavi stellam in linea recta a meridionali margine Insulæ Macræ (Posidonii) per septentrionalem partem Ponti Euxini, (medium maris Serenitatis) & M. Ætnam (Copernicum) producta: & linea a M. Sinai ad stellam fere stringebat littus sinus Sirbonidis. (M. Humorum).

F f

II. Tem-

II. Tempore observationis, (1.) stella erat in linea recta a lacu nigro majore (Platone) per partes orientales insulæ Cercinnæ (a Keplero versus ortum) producta.

III. Tempore observationis (2.) stella in linea per medium Paludis Mæotidis & medium M. Adriatici (per M. Crisium & S. medium) continuata.

IV. Tempore immersionis stellæ, sequentes lineæ rectæ ad illam coincidebant, & locum peripheriæ lunaris, ubi stella occultabatur, designabant.

1. A littore Pontus Euxini (M. Serenitatis) versus Cæciam spectante, per M. Ætnam (Copenicum).

2. A littore sinus Apollinis, per loca paludosa (sc. a littore S. Iridum, per Keplero).

3. A M. Sinai (Tychone) per littus meridionale S. Sirbonis (M. Humor).

V. Emergio stellæ contigit e regione M. Paropamisi, (Furnerii) & in linea recta a lacu nigro majore (Platone) per Byzantium (Menelaum) producta, quæ stringebat sinum extremum Ponti (M. Nectaris).

VI. Tempore observationis (7.) M. Porphyrites, (Aristarchus) margo septentrionalis L. Thespitis (Fracastorii) & stella in linea recta.

VII. Tempore observationis (8.) Lacus Hyperboreus superior (Hermes) medium Paludis Mæotidis (M. Crisii) & stella in linea recta.

*Occultatio Palilicii observata Vitembergæ Saxonum,*  
d. XXIII. Dec. St. N. A. CIO DCCXXXVIII. a Jo.  
Friderico Weidlero, R. S. Lond. S. &c.

Immersio	:	:	:	:	:	:	:	6.	27.	35.
Emersio	.	.	.	.	.	.	.	7.	29.	20.
Duratio	.	.	.	.	.	.	.	I.	I.	45.

# ANNOTATIONES.

1. Observatio a duobus spectatoribus simul perfecta; ego tubo astronomico IX pedum utebar, amicus & focius observationis per telescopium IV pedum lunam contuebatur.





2. Immersio & emersio facta in instanti: uno tamen prope minuto horario citius per longiorem, quam per breviorum tubum animadversa.

3. Appulit stella ad marginem lunæ orientalem, circa gradum 163. schematis lunæ plenæ mobilis Heveliani, Selenographiæ, p. 364. emergebat circa gradum 272. ejusdem schematis. Itaque linea recta, puncta immersionis & emersionis copulans, stringit extrema maris Humorū & Nubium & inter Pitatum & Mare Nubium transit.

Coelum tempore immersionis serenum non erat, sed tenues nubeculæ semper fere ante lunam & stellam oberrabant: ideoque figura stellæ oblonga, diu ante occultationem, per atmosphæræ vapores spectabatur.



Fig. 1.








Fig. 5.



TAB. II.



Mynd. Sulp.



Fig. 1.







Fig. 5.



Scala Minutorum Gradus.